




International Journal of Gerontology

journal homepage: <http://www.sgecm.org.tw/ijge/>



CME for 16.2

1. Which following symptom is NOT included in Fried's frailty phenotype?
(A) Weight loss
(B) Weakness (low handgrip strength)
(C) Exhaustion
(D) Depression
(E) Low physical activity
2. Compared with control group, which following statement about the patient navigators for transition care of heart failure is NOT true?
(A) Patient navigator group had higher cardiac rehabilitation enrollment.
(B) Patient navigator group had lower hospital-based utilization.
(C) Young patient (< 60 years old) in the patient navigator group had lower readmission rate.
(D) Patient navigator group had lower health care cost.
(E) Overall rate of readmission was similar in the two groups.
3. Reviewing the diagnostic test accuracy of physical examination tests in suspected patellofemoral osteoarthritis case, which test is most accurate?
(A) Crepitus sign
(B) Pain during activities
(C) Joint compression test
(D) Pain going up stairs
(E) Difficulty descending stairs
4. When elderly patients require surgical procedures on their heart valves and/or coronary arteries, which following parameter is associated with increased post-operative complications?
(A) Arterial hypertension
(B) Mini-mental State Examination
(C) Edmonton Frail Scale
(D) Society of Thoracic Surgeons (STS) scores
(E) All above
5. The probability of mortality increases in patients above 85 years of age who developed acute kidney injury in intensive care unit. Which following parameter has negative impact on the mortality?
(A) Increase in sequential organ failure assessment score
(B) Increase in ventilator workload
(C) Increase in serum lactate level
(D) Increase in body mass index
(E) Need for hemodialysis

Issue date: April 1st, 2022

Expiration date: July 31st, 2022

Volume 16 Issue 1 Answers:

1. (D)
2. (F)
3. (F)
4. (D)
5. (D)